

Catheterization and echocardiography confirmed the diagnosis of constrictive pericarditis which presumed to be due to old Tuberculosis infection .The patient underwent pericariectomy with good results.

Conclusion

Constrictive pericarditis should be suspected in patients with clinical features of right-sided heart failure.

Computed tomography (CT) is a useful method for accurate measurement of pericardial thickness and calcium distribution before pericardiectomy.

References

1. Michael Bergman , Janos Vitrai , Hertzel Salman. Constrictive pericarditis: A reminder of a not so rare disease . Eur J Intern Med 2006 Nov;17:457-64.
2. Little WC, Freeman GL . Pericardial disease. Circulation 2006, 113:1622-1632.

Ben Ahmed .Habib -Bouzouita .Khaled-Imen .Hamdi -Selmi .Khalifa -Boujnah .Mohamed Rachid

Department of Cardiology, Mongi Slim Hospital Tunis –Tunisia

Case report of a patient presenting both schizophrenia and Charcot-Marie-Tooth disease

Charcot-Marie-Tooth disease (CMT) is the most frequent inherited neuromuscular disorder (prevalence 17–40/100000) [1]. CMT comprises a group of disorders that affect peripheral nerves. CMT can be categorized into subtypes according to gene mutations [2]. Diagnosis is based on clinical findings and electromyography (EMG) and nervous conduction velocity characteristics.

CMT can rarely be associated with psychotic manifestations.

Psychosis is mainly caused by schizophrenia but can also be secondary to medical conditions or drug abuse.

Schizophrenia is the most frequent and debilitating psychotic disorder. It is the result of a complex group of genetic, psychological, and environmental factors.

Late adolescence and early adulthood are peak periods for the onset of schizophrenia. Its symptoms comprise the so-called positive symptoms or psychotic domain of schizophrenia. Negative symptoms of blunted affect, anhedonia, apathy, and alogia are a second symptom domain in schizophrenia.

Comorbidity between CMT and schizophrenia appears to be exceptional. This case report raises questions about the possible links between both conditions.

Case report

C. D. was diagnosed with CMT at the age of 16. He used to suffer from gait disturbances, while being an excellent student

with no symptoms suggestive of a psychotic disorder.

Clinical features included weakness of the foot and lower leg muscles, foot deformities: high arches and hammertoes. The clinical examination noticed absent achilles' tendon reflex and a superficial sensory disturbance. In addition to this peripheral neurogenic syndrome, a quadripyramidal syndrome was also noticed consisting of tendon hyperreflexia in four limbs (except the achilles' tendon reflex which was absent) and a positive bilateral Babinski sign.

An EMG was performed and showed a motor and sensory axonal neuropathy. Family tree suggested an autosomic dominant transmission.

The cerebro-medullar Magnetic Resonance Imaging was normal.

Thyroid function was normal.

HbS, HIV and syphilis were negative.

The patient was diagnosed with CMT associated with a pyramidal syndrome.

C.D. was diagnosed with schizophrenia in 2009, at the age of 27. The prodromal period lasted about one year consisting of deterioration of social functioning and social withdrawal.

The patient's clinical psychiatric features were: auditory hallucinations, persecutory delusion, behaviour disturbances consisting of self-neglect, obscene language and exhibitionism.

All other causes of psychosis were discussed and ruled out. The patient didn't take any medication that could account for his neurological nor psychiatric symptoms.

He was put on antipsychotics. Now he is on amisulpride 400mg qd and is well stabilized.

Family history was negative for schizophrenia.

Thus, C.D. had both schizophrenia and a form of CMT associated with a pyramidal syndrome. This involvement of the CNS in CMT does raise questions about the pathophysiology of both conditions and the possible links between them [3-5].

Conclusion

Comorbidity between schizophrenia and CMT is rare. It is even rarer and worth reporting when CMT with CNS involvement is concerned. It raises questions and suggests new theories about the pathophysiology of both diseases and the possible link between them [6]. No definite answer has been found so far. Additional research is needed to improve our knowledge of both diseases.

References

- 1) Ferrarin M, Bovi G, Rabuffetti M et al. Reliability of instrumented movement analysis as outcome measure in Charcot-Marie-Tooth disease: results from a multitask locomotor protocol. Gait and Posture 2011;34:36-43.
- 2) Verhoeven K, Claes K, Zuchner S et al. MFN2 mutation distribution and genotype/phenotype correlation in Charcot-Marie-Tooth type 2. Brain 2006;129:2093-102.
- 3) Thiam A, Sene Fd, Ndao A, Ndiaye M, Ndiaye I. Charcot-Marie-Tooth disease associated with a pyramidal syndrome: clinical, electrophysiological, and neuropathological study of neuro-muscular biopsies in 14 cases. Dakar Med 2002;47:182-7.
- 4) Barhoumi C, Amouri R, Ben Hamida C et al. Linkage of a new locus for autosomal recessive axonal form of Charcot-Marie-Tooth disease to chromosome 8q21.3. Neuromusc Disord 2001;11:27-34.
- 5) Barhoumi C, Machghoul S, Hintati F. Identification of a new genetic entity in the form of an autosomal dominant axonal Charcot-Marie-Tooth disease